

+ A new way to manage the incision and the outcome

Ensure your patients have the best post-op care to aid in the speed of recovery time and help improve scar appearance¹⁻⁴

Smith+Nephew

PICO[◇]
Single Use Negative Pressure
Wound Therapy System



The cascading consequences of SSIs

Financial costs



High-risk patients are likely to incur substantially **higher costs** with standard care for total joint procedures⁵


\$1,607
per case



Standard care for total joint procedures costs an additional **\$1,607 per case** due to the impact of Surgical Site Complications (SSCs)⁵



Patients with a **BMI ≥ 35** or an **ASA ≥ 3** can cost an additional **\$10,293 to \$11,296 per case**⁵



The consequence of an infection can lead to an additional **10 days before discharge**⁶

Personal costs

When patients experience complications they, their families and caregivers can all be impacted.⁷



Increased pain



Insecurity,
isolation



Loss of income



Inability to participate
in many daily activities,
social interactions


Know your patient. Know the risk factors.

All surgical procedures can be considered high-consequence. However, patients who have multiple risk factors are more likely to experience complications post-surgery. These outliers represent a small percentage but have a considerable impact on patient experience and outcomes.

Just one major risk factor or two or more moderate risk factors place patients at a high risk of surgical site complications.^{8,9}

Interrupt the pathway to complications

Uncontrolled edema is the root cause of many SSCs, which can reduce post-op mobility as persistent drainage/discharge can hamper recovery¹⁰



A preponderance of evidence shows that single-use NPWT improves clinical and economic outcomes for those at risk of SSCs; numerous orthopedic surgical departments have developed protocols for the prophylactic use of sNPWT.

The PICO[®] System helps provide advanced recovery

- The PICO Systems compressive forces help **improve perfusion, reducing lateral tension¹⁵⁻¹⁸ and dead space** that can lead to edema, while **increasing lymphatic drainage.¹⁹⁻²¹**
- When applied post-operatively clinicians and patients have noted **significant improvement, quality and appearance of scars.^{*1-4}**
 - The PICO System is indicated for open wounds and can be applied on dehisced wounds post-operatively, if needed. It can be applied in the acute, post-acute and physician office setting.

*vs standard dressings

Benefits of a unique mode of action with NPWT

PICO^o is the first NPWT System to be indicated to aid in the reduction of the incidence of both superficial and deep incisional SSIs for high-risk patients in Class I and II wounds, post-operative seroma, and dehiscence when used on closed surgical incisions.*

High moisture transmission rate²

to transpire as much as 80% of the wound fluid¹⁵

Silicone adhesive layer

protects the wound environment and helps to minimize trauma and pain on removal²²⁻²⁴

Super absorbent core

locks fluid away from wound²⁵


AIRLOCK^o Technology layer

distributes NPWT across the wound while enabling movement of fluid through the dressing²⁵



A wider zone of compression^{26,27}

Consistent delivery of negative pressure with PICO[®] sNPWT in clinical practice




Photographs courtesy of Dr. M. Brem, center photograph included in Nordmeyer M, et al.²⁸


Visible effects of PICO sNPWT on tissue surrounding the wound in patients who underwent spinal surgery (dressing boundary clearly visible on patient after removal)² *Individual results will vary*

Compression is dispersed across the entire tissue under the PICO sNPWT dressing^{26,27}

PICO sNPWT



tNPWT




CT scanning of tissue displacement under PICO sNPWT and tNPWT over 24hrs in a porcine tissue model.

Case studies

High-risk patient with total hip replacement

65-year-old female with hypertension, diabetes, BMI 35 kg/m², osteoarthritis



Individual results will vary

High-risk patient with knee implant

62-year-old male, medial compartment OA, Hyperlipidemia, Prostate CA



Individual results will vary

Preferred by patients, hospitals, surgeons

The PICO[◇] System's portable, canister-free design has been shown to increase patient satisfaction rates across the clinical spectrum.*²⁴

- Portable system allows patients the freedom to continue daily activities²⁹
- Waterproof dressing, allowing patients to shower²⁹
- Quiet system better enables patients to sleep²⁹
- Now offering therapy for up to 14 days with PICO 14 System

Reduce superficial SSCs by up to **76%** while also reducing exudate, length of stay and dressing changes⁶

Save an estimated **\$8,800** per high-risk patient following primary hip and knee arthroplasty, compared to standard care¹⁹



*vs tNPWT



Take your first step
to closure

Important Safety Information

The PICO[®] pumps contain a MAGNET. Keep the PICO pumps at least 4 inches (10 cm) away from other medical devices at all times. As with all electrical medical equipment, failure to maintain appropriate distance may disrupt the operation of nearby medical devices. For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for Use (IFU) prior to use. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Smith+Nephew representative or distributor if you have questions about the availability of Smith+Nephew products in your area.

Advanced Wound Management

Smith & Nephew, Inc.
Fort Worth, TX 76109
USA

Customer Care Center

T 800 876-1261
F 727 392-6914

◊Trademark of Smith+Nephew
All Trademarks acknowledged
© 2024 Smith & Nephew, Inc.

www.smith-nephew.com
www.possiblewithpico.com

PCME28-42080-0924

References: **1.** Tanaydin V, Beugels J, Andriessen A, Sawor JH, Van Der Hulst R. Randomized Controlled Study Comparing Disposable Negative-Pressure Wound Therapy with Standard Care in Bilateral Breast Reduction Mammoplasty Evaluating Surgical Site Complications and Scar Quality. *Aesthetic Plast Surg.* 2018;42(4):927-935. **2.** Sharp E. Single use NPWT for the treatment of complex orthopaedic surgical and trauma wounds; *Journal of Wound Care Cases Supplement* 2013. Vol 22. No 10. S5-9. **3.** Hudson DA, Adams KG, Van Huyssteen A, Martin R, Huddleston EM. Simplified negative pressure wound therapy: clinical evaluation of an ultraportable, no-canister system. *Int Wound J.* 2015;12(2):195-201. **4.** Smith & Nephew 2019. Pre-Clinical Assessment of Single-Use Negative Pressure Wound Therapy during in vivo Porcine Wound Healing. Internal Report. DS/19/313/R. **5.** Nherera LM, Trueman P, Karlakki SL. Cost-effectiveness analysis of single-use negative pressure wound therapy dressings (sNPWT) to reduce surgical site complications (SSC) in routine primary hip and knee replacements. *Wound Repair and Regeneration.* 2017 May; 25(3):474-82. **6.** Karlakki SL, Hamad AK, Whittall C, et al. Incisional negative pressure wound therapy dressings (iNPWTd) in routine primary hip and knee arthroplasties: A randomised controlled trial. *Bone Joint Res.* 2016;5(8):328-337. **7.** Andersson AE, Bergh I, Karlsson JU, Nilsson K. Patients' experiences of acquiring a deep surgical site infection: an interview study. *Am J Infect Control.* 2010 Nov; 38(9):711-7. **8.** Sugrue M, Ciprandi G, Djohan R, et al. World Union of Wound Healing Societies (WUWHS) Consensus Document. Closed surgical incision management: Understanding the role of NPWT. *Wounds Int* [Internet]. 2016. **9.** www.woundsinternational.com/wuwhs/view/consensus-document/closed-surgical-incision-management-understanding-the-role-of-npwt (Last accessed July 29, 2017). **10.** Ebert JR, Joss B, Jardine B, Wood DJ. Randomized trial investigating the efficacy of manual lymphatic drainage to improve early outcome after total knee arthroplasty. *Arch Phys Med Rehabil.* 2013 Nov;94(11):2103-11. doi: 10.1016/j.apmr.2013.06.009. Epub 2013 Jun 26. PMID: 23810354. **11.** Deakin AH, Iyayi-Igbinovia A, Love GJ. A comparison of outcomes in morbidly obese, obese and non-obese patients undergoing primary total knee and total hip arthroplasty. *Surgeon.* 2018 Feb;16(1):40-45. doi: 10.1016/j.surge.2016.10.005. Epub 2017 Jan 27. PMID: 28139371. **12.** Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six countries. *Journal of hospital infection.* Pages 1 -15. 2017. **13.** Chesney D, Sales J, Elton R, Brenkel JJ. Infection after knee arthroplasty a prospective study of 1509 cases. *J Arthroplasty.* 2008 Apr;23(3):355-9. doi: 10.1016/j.arth.2007.05.052. PMID: 18358372. **14.** Gristina AG. Biomaterial-centered infection: microbial adhesion versus tissue integration. *Science.* 1987 Sep 25;237(4822):1588-95. doi: 10.1126/science.3629258. PMID: 3629258. **15.** Malmsjo M, Huddleston E, Martin R. Biological effects of a disposable, canisterless negative pressure wound therapy system. *ePlasty.*2014;14:e15. **16.** Ma Z, Shou K, Li Z, et al. Negative pressure wound therapy promotes vessel destabilization and maturation at various stages of wound healing and thus influences wound prognosis. *Exp Ther Med.* 2016;11(4):1307-1317. **17.** Xia CY, Yu AX, Qi B, et al. Analysis of blood flow and local expression of angiogenesis associated growth factors in infected wounds treated with negative pressure wound therapy. *Mol Med Rep.* 2014;9(5):1749-1754. **18.** Loveluck J, Copeland T, Hill J, Hunt A, Martin R. Biomechanical modelling of the forces applied to closed incisions during single-use negative pressure wound therapy. *ePlasty.* 2016;16:e20. **19.** Kilpadi DV, Cunningham MR. Evaluation of closed incision management with negative pressure wound therapy (CIM): hematoma/seroma and involvement of the lymphatic system. *Wound Repair Regen.* 2011;19(5):588-596. **20.** Scalise A, Calamita R, Tartaglione C, et al. Improving wound healing and preventing surgical site complications of closed surgical incisions: a possible role of Incisional Negative Pressure Wound Therapy. A systematic review of the literature. *Int Wound J.* 2016;13(6):1260-1281. **21.** Glaser DA, Farnsworth CL, Varley ES, Nunn TA, Sayad-Shah M, Breisch EA, et al. Negative pressure therapy for closed spine incisions: a pilot study. *Wounds* 2012;24(11):308-16. **22.** Hudson DA, Adams KG, Van Huyssteen A, Martin R, Huddleston EM. Simplified negative pressure wound therapy: clinical evaluation of an ultraportable, no-canister system. *Int Wound J.* 2015;12(2):195-201. **23.** Payne C, Edwards D. Application of the Single Use Negative Pressure Wound Therapy Device (PICO) on a Heterogeneous Group of Surgical and Traumatic Wounds. *ePlasty.* 2014:152-166. **24.** Kirsner R, Cyaandi D, Reyzelamn A, Vayser D, Jaimes H. Randomized Controlled Trial on the Efficacy and Acceptance of a Single-use Negative Pressure Wound Therapy system versus Traditional Negative Pressure Wound Therapy in the Treatment of Lower Limb Chronic Ulcers (VLU and DFU). Paper presented at: EWMA; 2019; Gothenburg, Sweden. **25.** Smith+Nephew July 2018. PICO 7Y Non-NPWT Wound Model Summary. Internal Report. DS.18.260.R. **26.** Casey C. Consistent delivery of therapeutic negative pressure levels by a single use negative pressure wound therapy system (sNPWT)* in a wound model. Paper presented at: EWMA; 2019; Gothenburg, Sweden. **27.** Smith+Nephew 2021.PICO™ Pressure Mapping Study. Internal Report. DS/19/211/R - Part B. **28.** Nordmeyer M, Pauser J, Biber R, et al. Negative pressure wound therapy for seroma prevention and surgical incision treatment in spinal fracture care. *Int Wound J.* 2016;13(6):1176-1179. **29.** Hurd, T, Trueman, P, & Rossington, A. Use of portable, single use negative pressure wound therapy device in home care patients with low to moderately exuding wounds: a case series. *Ostomy Wound Management.* Volume 60, Issue 3. March 2014.